

STATE OF THE ART

Global Initiative for the Diagnosis, Management, and Prevention of Chronic Obstructive Lung Disease

The 2020 GOLD Science Committee Report on COVID-19 and Chronic Obstructive Pulmonary Disease

David M. G. Halpin¹, Gerard J. Criner^{2*}, Alberto Papi³, Dave Singh⁴, Antonio Anzueto^{5*}, Fernando J. Martinez^{6*}, Alvar A. Agusti⁷, and Claus F. Vogelmeier⁸; on behalf of the GOLD Science Committee

¹College of Medicine and Health, University of Exeter Medical School, University of Exeter, Exeter, United Kingdom; ²Lewis Katz School of Medicine, Temple University, Philadelphia, Pennsylvania; ³Department of Medical Sciences, Cardiorespiratory and Internal Medicine Unit, University of Ferrara, Ferrara, Italy; ⁴Manchester University NHS Foundation Trust, University of Manchester, Manchester, United Kingdom; ⁵Division of Pulmonary/Critical Care Medicine, South Texas Veterans Health Care System, University of Texas Health, University of Texas, San Antonio, Texas; ⁶Division of Pulmonary and Critical Care Medicine, Weill Cornell Medical College, New York, New York; ⁷Hospital Clinic, IDIBAPS, University of Barcelona, CIBERES, Barcelona, Spain; and ⁸Department of Medicine, Pulmonary, and Critical Care Medicine, the German Center for Lung Research, University Medical Center Giessen and Marburg, Philipps-University Marburg, Marburg, Germany

ORCID ID: 0000-0003-2009-4406 (D.M.G.H.).

Abstract

The severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) pandemic has raised many questions about the management of patients with chronic obstructive pulmonary disease (COPD) and whether modifications of their therapy are required. It has raised questions about recognizing and differentiating coronavirus disease (COVID-19) from COPD given the similarity of the symptoms. The Global Initiative for Chronic Obstructive Lung Disease (GOLD) Science Committee used established methods for literature review to present an overview of the management of patients with COPD during the COVID-19 pandemic. It is unclear whether patients with COPD are at increased risk of becoming infected with SARS-CoV-2. During periods of high community prevalence of COVID-19, spirometry should only be used when it is essential for COPD diagnosis and/or to assess lung function status for interventional procedures or surgery. Patients with COPD should follow basic infection control measures, including social distancing, hand washing, and wearing a mask or face covering. Patients should remain up to date with appropriate vaccinations, particularly annual influenza vaccination. Although data are limited, inhaled corticosteroids, long-acting bronchodilators, roflumilast, or chronic

macrolides should continue to be used as indicated for stable COPD management. Systemic steroids and antibiotics should be used in COPD exacerbations according to the usual indications. Differentiating symptoms of COVID-19 infection from chronic underlying symptoms or those of an acute COPD exacerbation may be challenging. If there is suspicion for COVID-19, testing for SARS-CoV-2 should be considered. Patients who developed moderate-to-severe COVID-19, including hospitalization and pneumonia, should be treated with evolving pharmacotherapeutic approaches as appropriate, including remdesivir, dexamethasone, and anticoagulation. Managing acute respiratory failure should include appropriate oxygen supplementation, prone positioning, noninvasive ventilation, and protective lung strategy in patients with COPD and severe acute respiratory distress syndrome. Patients who developed asymptomatic or mild COVID-19 should be followed with the usual COPD protocols. Patients who developed moderate or worse COVID-19 should be monitored more frequently and accurately than the usual patients with COPD, with particular attention to the need for oxygen therapy.

Keywords: chronic obstructive pulmonary disease; COVID-19; treatment; diagnosis

(Received in original form September 16, 2020; accepted in final form November 4, 2020)

†This article is open access and distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives License 4.0 (<http://creativecommons.org/licenses/by-nc-nd/4.0/>). For commercial usage and reprints, please contact Diane Gern (dgern@thoracic.org).

*G.J.C. and A.A. are Associate Editors and F.J.M. is Deputy Editor of *AJRCCM*. Their participation complies with American Thoracic Society requirements for recusal from review and decisions for authored works.

D.S. is supported by the National Institute for Health Research (NIHR) Manchester Biomedical Research Centre (BRC).

Author Contributions: D.M.G.H., G.J.C., A.P., and D.S. conducted the literature reviews and drafted the initial version of the report. All authors were involved in writing and critical review of subsequent versions of this manuscript and all approved the final version for submission for publication.

Correspondence and requests for reprints should be addressed to David M. G. Halpin, D.Phil., University of Exeter Medical School, College of Medicine and Health, University of Exeter, Exeter, UK, EX1 2LU. E-mail: d.halpin@nhs.net.

Am J Respir Crit Care Med Vol 203, Iss 1, pp 24–36, Jan 1, 2021

Copyright © 2021 by the American Thoracic Society

Originally Published in Press as DOI: 10.1164/rccm.202009-3533SO on November 4, 2020

Internet address: www.atsjournals.org

Contents

Risk of Infection with SARS-CoV-2

Investigations

Testing for SARS-CoV-2 Infection
Spirometry and Pulmonary
Function Testing
Bronchoscopy
Radiology

Protective Strategies for Patients with COPD

Differentiating COVID-19 Infection from the Daily Symptoms of COPD

Maintenance Pharmacological

Treatment for COPD during the COVID-19 Pandemic

Use of Nebulizers

Nonpharmacological Treatment for COPD during the COVID-19 Pandemic

Review of Patients with COPD during the COVID-19 Pandemic

Treatment of COVID-19 in Patients with COPD

Exacerbations of COPD

Systemic Corticosteroids

Antibiotics

Pulmonary and Extrapulmonary Complications

Anticoagulation

Ventilatory Support for Patients with COPD and COVID-19 Pneumonia

Rehabilitation

Follow-up of Patients with COPD Who Developed COVID-19

Conclusions

For patients with chronic obstructive pulmonary disease (COPD), the worry of developing coronavirus disease (COVID-19) as well as the effects of the pandemic on the basic functions of society and/or social services pertaining to their health imposes additional stressors to their condition. The COVID-19 pandemic has made routine management and diagnosis of COPD more difficult as a result of reductions in face-to-face consultations, difficulties in performing spirometry, and limitations in traditional pulmonary rehabilitation and home care programs. Patients have also faced shortages of medication (1).

The dramatic spread of the SARS-CoV-2 virus has been accompanied by an enormous number of publications on the virus and its consequences. The statements made in this report (Table 1) use the published Global Initiative for Chronic Obstructive Lung Disease (GOLD) approach to data review and should be seen as *provisional* based on the best assessment of the current evidence, including papers published or available up to the September 11, 2020. As new evidence emerges, this report will be updated at regular intervals, and updates will be posted on the GOLD website (www.goldcopd.org).

Risk of Infection with SARS-CoV-2

It appears that the spike protein of the virus binds to ACE2 (angiotensin-converting enzyme 2) during viral attachment to host cells and that viral entry is also facilitated by TMPRSS2 (transmembrane serine protease 2) (2). Differences in the expression of ACE2 and TMPRSS2 may modulate the individual susceptibility to and clinical course of SARS-CoV-2 infection. ACE2

mRNA expression is increased in COPD (3, 4) and may be modulated by inhaled corticosteroid (ICS) use (3, 5).

It is not known definitively yet whether having COPD affects the risk of becoming infected with SARS-CoV-2. Very few population studies using random sampling have assessed risk factors for testing positive for SARS-CoV-2, most have looked at samples of patients referred for testing or presenting with symptoms, and very few contain information on comorbidities. A population survey with random sampling found no increased risk of infection (6). Similarly, most studies of people in the community tested for SARS-CoV-2 have not shown chronic respiratory disease as an independent risk factor for testing positive (7, 8), although at least one has (9).

Many studies reporting the comorbidities of patients admitted to the hospital with COVID-19 have suggested a lower prevalence of COPD than would be expected from population prevalence (10, 11); these findings are limited by small sample sizes and incomplete data on comorbidities. A large study with comprehensive data on comorbidities showed a high prevalence of COPD among those admitted (19%) (12), although many patients had multiple comorbidities, and a further study of a primary care cohort of 8.28 million patients showed having COPD was an independent risk factor for hospital admission (hazard ratio, 1.55; 95% confidence interval, 1.46–1.64) (9).

COPD has also been reported to independently increase the risk of severe disease or death in some series (12–15) but not all (9, 16, 17). Many factors have been proposed to account for the increased risk for poor outcomes, including prior poor adherence to therapy, difficulties performing self-management, limited access to care

during the pandemic, and a reduced pulmonary reserve (18, 19). There is evidence of a fall in hospitalization rates for COPD during the pandemic (20, 21). The reasons for this remain unclear, but patients experiencing symptoms of an exacerbation should be evaluated in the usual way during the pandemic and hospitalized if necessary.

There are currently no peer-reviewed studies that have evaluated the effect of smoking on the risk of infection with SARS-CoV-2, but studies suggest that smoking is associated with increased severity of disease and risk of death in hospitalized patients with COVID-19 (22).

In summary, on current evidence, patients with COPD do not seem to be at a greatly increased risk of infection with SARS-CoV-2, but this may reflect the effect of protective strategies. They are at an increased risk of hospitalization for COVID-19 and may be at increased risk of developing severe disease and death.

Investigations

Testing for SARS-CoV-2 Infection

Patients with COPD presenting with respiratory symptoms, fever, or other symptoms suggesting SARS-CoV-2 infection, even if mild, should be tested for possible infection (Figure 1). False-negative RT-PCR tests have been reported in patients with computed tomographic (CT) scan findings of COVID-19, who eventually tested positive with serial sampling (23). If patients with COPD have been exposed to someone with a known COVID-19 infection, they should contact their healthcare provider to define the need for specific testing. Antibody testing may be used to support clinical assessment of patients who present late.

Detection of SARS-CoV-2 does not exclude the potential for coinfection with other respiratory pathogens (24). The U.S. CDC encourages testing for other causes of respiratory illness in addition to testing for SARS-CoV-2 depending on patient age, season, or clinical setting.

Some patients experience reactivation of long-lasting virus carriage or become reinfected, and this might be influenced by comorbidities or drugs that hamper the immune response (25). Repeat testing should be performed in patients with suspected recurrence or relapse of COVID-19.

The lung microbiome is different in patients with COPD versus healthy subjects (26). The lung microbiome can modify the immune response to viral infections but, to date, there is no direct evidence from human or animal studies on the role of lung microbiome in modifying COVID-19 disease (27) nor on its potential effects in patients with COPD.

Spirometry and Pulmonary Function Testing

Performing spirometry and pulmonary function testing may lead to SARS-CoV-2 transmission as a result of coughing and droplet formation during the tests (28, 29). During periods of high prevalence of COVID-19 in the community, spirometry should be restricted to patients requiring urgent or essential tests for the diagnosis of COPD and/or to assess lung function status for interventional procedures or surgery. The American Thoracic Society and European Respiratory Society have provided recommendations regarding testing and precautions that should be taken (28, 29). Whenever possible, patients should have an RT-PCR test for SARS-CoV-2 performed, and the results should be available before performing the test. Patients with a positive RT-PCR test should normally have the test delayed until negative.

When routine spirometry is not available, home measurement of peak expiratory flow combined with validated patient questionnaires could be used to support or refute a possible diagnosis of COPD (30–33). However, peak expiratory flow does not correlate well with the results of spirometry (34–36), has low specificity (37), and cannot differentiate obstructive and restrictive lung function abnormalities. When making a diagnosis of COPD, airflow obstruction could also be confirmed by giving patients a personal electronic portable

spirometer (38, 39), instructing them in their use, and observing them in their homes using video conferencing technology.

Bronchoscopy

In some patients with COPD, diagnostic and therapeutic bronchoscopy may be required during the COVID-19 pandemic. An elective bronchoscopy should be delayed until patients have a negative PCR test (40, 41). In urgent cases in which COVID-19 infection status is unknown, all cases should be managed as if positive. A disposable bronchoscope should be used if available (40), and staff should wear personal protective equipment.

Radiology

Chest radiography is insensitive in mild or early COVID-19 infection (42) and is not routinely indicated as a screening test for COVID-19 in asymptomatic individuals. Chest radiography is indicated in patients with COPD who have moderate-to-severe symptoms of COVID-19 and for those with evidence of worsening respiratory status (43) (Figure 1). COVID-19 pneumonia changes are mostly bilateral (44). Chest radiography can be useful for excluding or confirming alternative diagnoses (e.g., lobar pneumonia, pneumothorax, or pleural effusion). Point-of-care lung ultrasound can also be used to detect the pulmonary manifestations of COVID-19 (45).

CT screening may show evidence of pneumonia in asymptomatic individuals infected with SARS-CoV-2 (46), and false-negative RT-PCR tests have been reported in patients with CT findings of COVID-19 who eventually tested positive (23).

Recommendations have been made on the use of CT scans as part of diagnostic testing and severity assessment in COVID-19 (43), and there are no special considerations for patients with COPD. The initial features of COVID-19 on CT scans and their progression over time have been reviewed (47). Patients with COPD and COVID-19 have an increased prevalence of ground-glass opacities, local patchy shadowing, and interstitial abnormalities on CT scans than patients without COPD (48). A small case series of patients with emphysema and COVID-19 found that many had bilateral ground-glass opacities with areas of consolidation; however, the pattern was variable, and patients had more pronounced disease in the lung bases (49).

The availability of CT scans may be limited by infection-control requirements (50), and where access to CT scans are limited, chest radiography may be preferred for patients with COVID-19 unless features of respiratory worsening warrant the use of CT scans. An increased occurrence of deep venous thrombosis and pulmonary thromboembolism has been reported in patients with COVID-19 (51–56); if pulmonary embolism is suspected, chest CT angiography should be performed.

Protective Strategies for Patients with COPD

Patients with COPD should follow basic infection-control measures to help prevent SARS-CoV-2 infection, including social distancing and washing hands (Table 2). Wearing a mask or face covering can reduce

Table 1. Key Points

<p>Patients with COPD presenting with new or worsening respiratory symptoms, fever, and/or any other symptoms that could be COVID-19 related, even if these are mild, should be tested for possible infection with SARS-CoV-2.</p> <p>Patients should keep taking their oral and inhaled respiratory medications for COPD as directed, as there is no evidence that COPD medications should be changed during the COVID-19 pandemic.</p> <p>During periods of high prevalence of COVID-19 in the community, spirometry should be restricted to patients requiring urgent or essential tests for the diagnosis of COPD and/or to assess lung function status for interventional procedures or surgery.</p> <p>Physical distancing and shielding, or sheltering-in-place, should not lead to social isolation and inactivity. Patients should stay in contact with their friends and families by telecommunication and continue to keep active. They should also ensure they have enough medication.</p> <p>Patients should be encouraged to use reputable resources for medical information regarding COVID-19 and its management.</p>

Definition of abbreviations: COPD = chronic obstructive pulmonary disease; COVID-19 = coronavirus disease; SARS-CoV-2 = severe acute respiratory syndrome coronaviruses 2.

the risk of *spreading* infection (source control) (57). The efficacy of masks and respirators in *protecting* patients against infection is unknown, but both surgical masks and N95 respirators were effective in preventing influenza-like illness and laboratory-confirmed influenza among healthcare workers (58). The American College of Chest Physicians, American Lung Association, American Thoracic Society, and the COPD Foundation have issued a joint statement on the importance of patients with chronic lung disease wearing facial coverings during the COVID-19 pandemic (59).

Wearing a tight-fitting N95 mask introduces an additional inspiratory resistance. Respiratory rate, peripheral oxygen saturation, and exhaled CO₂ levels were adversely affected in patients with COPD wearing an N95 mask for 10 minutes at rest followed by 6 minutes of walking (60); however, wearing a surgical mask does not appear to affect ventilation even in patients with severe airflow limitation (61). In some countries where wearing face masks is compulsory in certain settings, exemptions can be made for patients who are breathless and cannot

tolerate wearing a mask. Whenever possible patients should wear masks. In most cases, a looser face covering or even a face shield may be tolerable and effective (62, 63).

The normal rules for patients on long-term oxygen therapy should be followed if air travel is planned (64, 65), although patients should avoid travel unless essential. Supplementary oxygen should be delivered by nasal cannula (66), with a surgical mask worn and distancing maintained.

Shielding, or sheltering-in-place, is a way to protect people who are extremely vulnerable from coming into contact with coronavirus. It is an alternative to full-scale physical distancing measures or lockdowns. It has been introduced in some countries for patients with severe COPD. In the United Kingdom, patients with COPD were advised to shield if they had an FEV₁ < 50%, an mMRC ≥ 3, a history of hospitalization for an exacerbation, or required long-term oxygen therapy or noninvasive ventilation (NIV). Modeling suggests shielding is an effective strategy to protect individuals and control the impact of SARS-CoV-2 (67). If patients with COPD are asked to shield, it is important that they

are given advice about keeping active and exercising as much as possible while shielded. Plans should be made to ensure supplies of food, medications, oxygen, supportive health services, and other basic necessities can be maintained.

There are likely to be particular challenges in using shielding in low- and middle-income countries, including the fact that many families will not be able to designate a separate room for high-risk individuals and may rely on the income or domestic support that these individuals provide (68).

Differentiating COVID-19 Infection from the Daily Symptoms of COPD

Differentiating the symptoms of COVID-19 infection from the usual symptoms of COPD can be challenging. Cough and breathlessness are found in over 60% of patients with COVID-19 but are usually also accompanied by fever (over 60% of patients) as well as fatigue, confusion, diarrhea, nausea, vomiting, muscle aches and pains, anosmia, dysgeusia, and headaches (12).




Figure 1. Chronic obstructive pulmonary disease and severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection: clinical features, abnormal investigations, and possible interventions at different stages of the disease. ARDS = acute respiratory distress syndrome; BNP = brain natriuretic peptide; COPD = chronic obstructive pulmonary disease; COVID-19 = coronavirus disease; CRP = C-reactive protein; CT = computed tomography; CXR = chest radiograph; HFNT = high-flow nasal therapy; IMV = invasive mechanical ventilation; LDH = lactate dehydrogenase; NIV = noninvasive ventilation; PCT = procalcitonin; PFT = pulmonary function tests; PR = pulmonary rehabilitation; SIRS = systemic inflammatory response syndrome; SOB = shortness of breath; SpO₂ = peripheral oxygen saturation; VTE = venous thromboembolism.

Table 2. Key Points for the Management of Stable Chronic Obstructive Pulmonary Disease during the COVID-19 Pandemic

Protective strategies
Follow basic infection control measures
Wear a face covering
Consider shielding/sheltering in place
Investigations
Only essential spirometry
Pharmacotherapy
Ensure adequate supplies of medications
Continue unchanged including ICS
Nonpharmacological therapy
Ensure annual influenza vaccination
Maintain physical activity

Definition of abbreviations: COVID-19 = coronavirus disease; ICS = inhaled corticosteroids.

In COVID-19, symptoms may be mild at first, but rapid deterioration in lung function may occur (Figure 1). The prodrome of milder symptoms is especially problematic in patients with underlying COPD who may already have diminished lung reserve. The lack of recognition of the prodromal symptoms may delay early diagnosis, and preliminary data suggest that patients with COPD reporting exacerbations and suspected of having COVID-19 infection were infrequently tested for its presence (69). A high index of suspicion for COVID-19 needs to be maintained in patients with COPD who present with symptoms of an exacerbation, especially if accompanied by fever, impaired taste or smell, or gastrointestinal complaints.

Persistent symptoms in patients with COPD may cause diagnostic difficulty. A recent study found that only 65% of people had returned to their previous level of health 14–21 days after testing positive for SARS-CoV-2 (70). Some patients continued to experience cough, fatigue, and breathlessness for weeks and a smaller proportion for months (70–72). Delayed recovery was more common in people with multiple chronic medical conditions but was not specifically linked to having COPD (70).

Maintenance Pharmacological Treatment for COPD during the COVID-19 Pandemic

The use of inhaled and systemic corticosteroids has been controversial in the prevention and treatment of COPD during the COVID-19 pandemic. ICS have an overall protective effect against exacerbations in patients with COPD

and a history of exacerbations (64). However, there is an increased risk of pneumonia associated with ICS use, raising concerns that immunosuppression with ICS could increase susceptibility to infections in some individuals.

Laboratory experiments show that corticosteroids reduce the production of antiviral IFNs (type I and III), increasing the replication of the rhinovirus and the influenza virus (73–75). In contrast, other laboratory data show that corticosteroids and long-acting bronchodilators can reduce the replication of coronaviruses, including SARS-CoV-2 (76). These laboratory experiments suggesting a potential protective effect of ICS against COVID-19 have not been validated by clinical studies.

A systematic literature review identified no clinical studies in patients with COPD concerning the relationship between ICS use and clinical outcomes with coronavirus infections, including COVID-19, SARS, and Middle East Respiratory Syndrome (MERS) (77). A more recent study has shown ICS use in COPD was not protective and raised the possibility that it increased the risk of developing COVID-19 (78), but the results are likely to be confounded by the indication for ICS (79). There are no conclusive data to support the alteration of maintenance COPD pharmacological treatment either to reduce the risk of developing COVID-19 or, conversely, because of concerns that pharmacological treatment may increase the risk of developing COVID-19.

Similarly, there are no data on the use of long-acting bronchodilators, roflumilast, or macrolides in patients with COPD and clinical outcomes or risk of SARS-CoV-2 infection; thus, unless evidence emerges, these patients should continue these medications required for COPD.

Use of Nebulizers

Aerosol therapy increases the droplet generation and risk of disease transmission. Although most of the aerosol emitted comes from the device (80, 81), there is a risk that patients may exhale contaminated aerosol and droplets produced by coughing when using a nebulizer and may be dispersed more widely by the driving gas. SARS-CoV-2 has been shown to be viable in aerosols for up to 3 hours (82), and transmission to healthcare workers exposed to a hospitalized patient with COVID-19 receiving nebulized therapy has been reported (83). If possible, pressurized

metered-dose inhalers, dry powder inhalers, and soft mist inhalers should be used for drug delivery instead of nebulizers. The risks of nebulized therapy spreading infection to other people in patient's homes can be minimized by avoiding use in the presence of other people and ensuring that the nebulizer is used near open windows or in areas of increased air circulation (84).

Nebulizers may be needed in patients who are critically ill with COVID-19 receiving ventilatory support. In this case, it is vital to keep the circuit intact and prevent the transmission of the virus. Using a mesh nebulizer in patients who are ventilated allows for the addition of medication without requiring the circuit to be broken for aerosol drug delivery (85).

Nonpharmacological Treatment for COPD during the COVID-19 Pandemic

During the COVID-19 pandemic, patients with COPD should continue with their nonpharmacological therapy (64). Patients should receive their annual influenza vaccination, although the logistics of providing these while maintaining social distancing will be challenging (86). There is no reason to modify palliative care approaches because of COVID-19.

Many pulmonary rehabilitation programs have been suspended during the pandemic to reduce risks of spreading SARS-CoV-2. When case rates are high, center-based rehabilitation is not appropriate. Patients should be encouraged to keep active at home and can be supported by home-based rehabilitation programs, which, although likely to be less effective than traditional pulmonary rehabilitation with supervision (64), are likely to be better than not offering rehabilitation. Technology-based solutions, such as web-based or smartphone applications (87), may be useful to support home rehabilitation during the pandemic. As programs are restarted, general principles of infection control should be applied and local guidance followed (88).

Review of Patients with COPD during the COVID-19 Pandemic

To minimize the spread of SARS-CoV-2, many health systems have reduced

Table 3. Key Points for the Management of Patients with Chronic Obstructive Pulmonary Disease and Suspected or Proven COVID-19

SARS-CoV-2 testing	
	Swab/saliva PCR if new or worsening respiratory symptoms, fever, and/or any other symptoms that could be COVID-19 related
Other investigations	
	Avoid spirometry unless essential
	Consider CT for COVID-19 pneumonia and to exclude other diagnoses (e.g., PE)
	Avoid bronchoscopy unless essential
	Assess for coinfection
COPD pharmacotherapy	
	Ensure adequate supplies of medication
	Continue maintenance therapy unchanged including ICS
	Use antibiotics and oral steroids in line with recommendations for exacerbations
	Avoid nebulization when possible
COPD nonpharmacological therapy	
	Maintain physical activity as able
Protective strategies	
	Follow basic infection control measures
	Maintain physical distancing
	Wear a face covering
COVID-19 therapy	
	Use systemic steroids and remdesivir as recommended for patients with COVID-19
	Use HFNT or NIV for respiratory failure if possible
	Use invasive mechanical ventilation if HFNT or NIV fails
	Post COVID-19 rehabilitation
	Ensure appropriate post-COVID-19 follow up

Definition of abbreviations: COPD = chronic obstructive pulmonary disease; COVID-19 = coronavirus disease; CT = computed tomography; HFNT = high-flow nasal therapy; ICS = inhaled corticosteroids; NIV = noninvasive ventilation; PE = pulmonary embolism; SARS-CoV-2 = severe acute respiratory syndrome coronaviruses 2.

face-to-face visits and introduced remote consultations using online, phone, and video links. Routine review of patients with COPD can be undertaken remotely (89), and GOLD has produced a tool to support these interactions that includes instructions on how to prepare for the remote visit and set the visit agenda with the patient, and it provides a standardized checklist for follow up (www.goldcopd.org).

Treatment of COVID-19 in Patients with COPD

Randomized clinical trials of treatments targeting COVID-19 have focused on antiviral agents and antiinflammatory treatments. Some have produced positive results, including the antiviral drug remdesivir (90) and systemic steroids for hospitalized patients with severe COVID-19 (91). Subgroup analysis on COPD patients has not been presented in these trials.

In the absence of subgroup data, we recommend that patients with COPD and COVID-19 should be treated with the same standard of care treatments as other patients with COVID-19 (Figure 1). Importantly,

there are no known drug interactions between remdesivir and inhaled COPD treatments. Furthermore, we advocate that patients with COPD should be included in randomized controlled trials of COVID-19 treatments and that subgroup analyses of their outcomes are presented.

Exacerbations of COPD

The prevention and treatment of exacerbations are important goals in COPD management (64). COVID-19 infection has introduced unique obstacles to the prevention and management of exacerbations (19). These include limited access to therapies because of their use for patients with COVID-19 without COPD, disruptions in global supply chains, and the inability of patients to afford medications owing to economic hardships associated with the pandemic (19). Conversely, as countries went into lockdown and industrial activities shut down, pollutant emissions reduced substantially and environmental air quality improved (92). This could have contributed to the reported reductions of in-hospital admissions for

COPD during the COVID-19 pandemic (20, 21)

Coronaviruses are among the respiratory viruses that trigger COPD exacerbations (93). To date, MERS-CoV, SARS-CoV, and SARS-CoV-2 infection have not been reported in COPD exacerbations. Nonetheless, patients with COPD and SARS-CoV-2 infection presenting with respiratory symptoms requiring changes in their maintenance medications would fulfill the definition of an exacerbation (64). Distinguishing the symptoms of a typical exacerbation from COVID-19 infection can be extremely difficult, as many of the symptoms overlap. If a COVID-19 infection is suspected, then RT-PCR testing should be conducted (Table 3). If COVID-19 infection is confirmed, then treatment for COVID-19 infection should be conducted regardless of the presence of COPD.

SARS-CoV-2 infection causes a distinct pattern of pathophysiological changes, including vascular injury, pneumonitis associated with hypoxemia, coagulopathy, high levels of systemic inflammation (“cytokine storm”), and multiorgan involvement (94, 95). These features are very different from typical COPD exacerbations (96). However, SARS-CoV-2 infection may resemble an exacerbation of COPD. Fever, anorexia, myalgias, and gastrointestinal symptoms are more frequently reported in COVID-19 than in exacerbations of COPD, whereas sputum production occurs in both. Pronounced lymphopenia is a common finding of SARS-CoV-2 infection (53, 97). Patients with COPD who develop COVID-19 reported more severe fatigue, dyspnea, and diarrhea than those without COPD (48).

In patients with COVID-19, lymphopenia, thrombocytopenia, elevated D-dimer, C-reactive peptide, procalcitonin, creatinine kinase, transaminases, creatinine, and lactate dehydrogenase are independently associated with higher risk of poor outcomes (98). There is no reason to suspect that this is different in patients with COPD and COVID-19 (Figure 1).

Systemic Corticosteroids

Caution has been raised about the widespread use of systemic corticosteroids in patients with COVID-19 (99, 100). Observational studies in patients with SARS and MERS reported no association between

systemic corticosteroids (often at high dose) and improved survival, but they suggested that corticosteroids induced side effects, including osteonecrosis, and reduced viral clearance (101–104). The World Health Organization initially recommended against the routine use of corticosteroids in COVID-19 infection at the beginning of the pandemic, except in two clinical settings: acute respiratory distress syndrome (ARDS) and COPD exacerbations, in which specific indications for systemic corticosteroids were recognized (105).

A large, randomized trial in hospitalized patients with COVID-19 has shown that dexamethasone treatment at 6 mg/d for up to 10 days reduced mortality in patients receiving either invasive mechanical ventilation (IMV) or oxygen alone (91). A small observational study has also reported that methylprednisolone use was associated with improved survival in patients with COVID-19 and ARDS (106). Further studies have also reported the benefits of systemic glucocorticoids on reduction of mortality at 28 days in patients with COVID-19 pneumonia, especially those that are not on IMV or on pressor support (107).

Systemic steroids should be used in COPD exacerbations according to the usual indications (64) whether or not there is evidence of a SARS-CoV-2 infection, as there is no evidence that this approach modifies the susceptibility to a SARS-CoV-2 infection or worsens outcomes (Figure 1).

Antibiotics

Antibiotic treatment for a COPD exacerbation is indicated if patients have at least two of the three cardinal symptoms, including increased sputum purulence, or if the patient requires mechanical ventilation (64).

Bacterial coinfections have been reported infrequently in COVID-19 (108). However, the risk of coinfections increases with the severity of COVID-19. Bacterial coinfections have been detected by multiplex PCR testing in up to 46% of samples collected in a small cohort of patients with COVID-19 admitted to an ICU (109). Diagnosing coinfection in patients with COVID-19 may be difficult, particularly in critically ill subjects, as the clinical presentation, biomarkers, and imaging data may be unhelpful. In practice, most hospitalized patients, particularly the

severe ones, have been prescribed empirical antibiotic therapy (97, 110). Current World Health Organization guidelines recommend broad-spectrum antibiotics in patients with severe COVID-19, guided by the local/national guidelines and in milder COVID-19 infections when there is clinical suspicion of a bacterial infection (105). In the absence of specific studies, these general considerations would also apply to patients with COPD infected with SARS-CoV-2.

Antibiotics should be used in COPD exacerbations according to the usual indications (64) whether or not there is evidence of a SARS-CoV-2 infection, particularly as patients with COPD who develop COVID-19 are reported, to more frequently develop bacterial or fungal coinfections (48).

Pulmonary and Extrapulmonary Complications

ARDS may be part of COVID-19 and could be considered the major pulmonary complication of COVID-19 (111), with viral infection in areas of ongoing active injury contributing to persistent and temporally heterogeneous lung damage (112). Some early reports suggested that ARDS in this setting may differ from the typical ARDS (113, 114). Subsequent studies, however, suggested that classical ARDS also presents with a large variation in lung severity (115), and there is considerable overlap between patients with classical ARDS and patients with COVID-19 (116, 117). Whether the long-term consequences of this form of ARDS differs from the fibrotic lesions described previously is unclear (118, 119).

Although the respiratory tract is the main target of COVID-19, extrapulmonary involvement is frequent and contributes to morbidity, disability, and mortality (95, 120). Renal, cardiac, nervous, cutaneous, hepatic, and gastrointestinal manifestations occur (121). It remains unclear, however, if these manifestations are directly caused by infection of SARS-CoV-2 or secondary phenomena, including inappropriate or overwhelming immune responses, angiopathy, treatment, or ischemic damages because of the impairment of the respiratory functions. Concomitant respiratory comorbidities, such as COPD, may aggravate these processes. Compared with lung viral

load, lower levels of SARS-CoV-2 have been reported in the kidneys, liver, heart, and brain (122), suggesting secondary rather than primary involvement of these organs.

Anticoagulation

COVID-19 has been associated with a hypercoagulable state (51), and venous thromboembolism rates in both ICU and ward patients are two- to fourfold higher than expected despite thromboprophylaxis with a low-molecular-weight heparin (LMWH) or unfractionated heparin (123). Patients with COPD are already at an increased risk for venous thromboembolism (124, 125), and those hospitalized with COVID-19 should receive pharmacologic thromboprophylaxis (Figure 1). In response to the high rates despite prophylactics, many institutional protocols have adopted intermediate-intensity (i.e., twice-daily LMWH rather than once daily) or even a therapeutic-intensity dose strategy for thromboprophylaxis (126). Generally, LMWH is favored over unfractionated heparin to reduce staff exposure, but clinicians should follow local guidelines on the dosing and drug.

Ventilatory Support for Patients with COPD and COVID-19 Pneumonia

The prevalence of hypoxic respiratory failure in patients with COVID-19 is around 19% (127). Ventilatory support has been used in up to 20% of patients who develop severe hypoxemia because of COVID-19 (128), and approximately 5% of patients require ICU care and advanced respiratory support (129). Patients requiring ventilatory support have a high risk of mortality (14, 130), and COPD has been reported to increase the risk of respiratory failure and ICU admissions in some but not all studies (9, 13).

There is a wide variation (2.3–33%) in the reported rates of use of IMV in hospitalized patients with moderate-to-severe hypoxic respiratory failure because of COVID-19 (131). This may, in part, reflect differences in the use of NIV and high-flow nasal therapy (HFNT) (131), possibly as a result of advocacy of early intubation during the pandemic's initial phases because of concerns about viral dissemination (132, 133). Data supporting those concerns are lacking (134).

Although early reports showed mixed outcomes (135), several studies have now shown that HFNT significantly reduces rates of intubation and IMV, although with variable effects on mortality (136, 137). HFNT should be considered in preference to NIV for acute hypoxemic respiratory failure despite conventional oxygen therapy, as it may have a lower failure rate (138–140). Prone positioning has also been suggested for awake nonintubated patients with hypoxemia (141).

NIV is the normal standard of care for patients with COPD and acute respiratory failure (64). NIV may be beneficial for the treatment of hypercapnic respiratory in patients with COPD and COVID-19 pneumonia, but it also has the potential to worsen lung injury as a result of high transpulmonary pressures and V_T (142). Patients on HFNT or NIV should be monitored closely for worsening, and early intubation and IMV with adoption of a protective lung strategy, similar to that used in other forms of ARDS, should be considered (143, 144). A PaO_2/FiO_2 ratio < 150 mm Hg may be a useful indicator for NIV failure and increased risk of mortality (145).

Evidence on the effects of extracorporeal membrane oxygenation in COVID-19 is scant and retrospective (131, 146–150). The indications in COVID-19 are similar to the indications for other causes of ARDS (151, 152), and extracorporeal membrane oxygenation should be considered only after other strategies fail to achieve goals of oxygenation or ventilation (147, 148, 150).

Aerosol generation can occur when any form of additional pressures or flows are applied to the upper or lower respiratory tract (153). Data regarding aerosol dispersion with the use of NIV are limited and contradictory (81, 153–155); however, staff should use appropriate personal protective equipment (140, 156) and viral filters fitted to exhalation ports of invasive or noninvasive ventilation devices. Isolation hoods have also been suggested by some to be used to further decrease staff exposure (157).

Rehabilitation

Patients with COPD and COVID-19 are particularly at risk for poor nutritional status and skeletal muscle loss (158). Hospital treatment should therefore include dietary support and early mobilization. Mechanical ventilation, sedation, and prolonged bed rest may lead to posttraumatic stress disorder (159) and respiratory, cognitive, and mental health impairments as well as physical deconditioning (160, 161). Older people and patients with COPD are more susceptible to these consequences (162, 163).

Rehabilitation should be provided to all patients with COPD and COVID-19, particularly to those that have been more severely affected or required ICU admission. A multinational task force has recommended early rehabilitation during the hospital admission and the screening for traits treatable with rehabilitation in all patients at discharge and at 6–8 weeks after discharge for patients with severe COVID-19 (164).

Follow-up of Patients with COPD Who Developed COVID-19

Approximately 30% of patients with SARS or MERS had persistent lung abnormalities and abnormal radiology that were consistent with fibrotic lung disease after their acute illness (165, 166). There are not yet long-term studies on the follow-up of patients with COVID-19, nor recommendations for monitoring these patients (143, 167), thus the follow up of patients with COPD who developed COVID-19 is still based on expert opinion and consensus. The intensity of the monitoring obviously depends on the severity of the COVID-19 episode.

Patients who developed mild COVID-19 should be followed with the usual protocols used for patients with COPD (64). Patients who developed moderate COVID-19, including hospitalization and pneumonia but no

respiratory failure, should be monitored more frequently and accurately than the usual patients with COPD, with particular attention to the need for oxygen therapy.

If chest X-ray abnormalities have not resolved at hospital discharge, a chest X-ray and possibly a CT scan should be considered at 6 months to 1 year. Complications occurring during/after the COVID-19 episode should also be monitored.

Patients with COPD are at higher risk of developing severe COVID-19 (143, 168), and multimorbid survivors frequently have required prolonged ICU stays (143). Until we have evidence from prospective studies, survivors of severe COVID-19 with COPD should be considered at a high risk of developing a “critical illness” (169) or a “chronic critical illness” (170), a severe heterogeneous condition linked not only to the acute infectious episode but also to the underlying conditions before they became severely ill (161).

There are informative candidate models for the comprehensive management of complex care delivery that are already published and undergoing study in a primary care setting, and these may be adapted for application after COVID-19 (171).

Conclusions

There is little direct evidence about management of COVID-19 in people with COPD. Clinicians should maintain a high level of suspicion of COVID-19 in patients with COPD presenting with new or worsening respiratory symptoms, fever, and/or any other symptoms that could be COVID-19 related, even if these are mild, and should test for SARS-CoV-2. Patients should keep taking their oral and inhaled respiratory medications for COPD as directed, as there is no evidence that COPD medications should be changed during this COVID-19 pandemic. ■

Author disclosures are available with the text of this article at www.atsjournals.org.

References

- Mahase E. Covid-19: increased demand for steroid inhalers causes “distressing” shortages. *BMJ* 2020;369:m1393.
- Hoffmann M, Kleine-Weber H, Schroeder S, Kruger N, Herrler T, Erichsen S, et al. SARS-CoV-2 cell entry depends on ACE2 and TMPRSS2 and is blocked by a clinically proven protease inhibitor. *Cell* 2020;181:271–280, e8.
- Maes T, Bracke K, Brusselle GG. COVID-19, asthma, and inhaled corticosteroids: another beneficial effect of inhaled corticosteroids? *Am J Respir Crit Care Med* 2020;202:8–10.
- Leung JM, Yang CX, Tam A, Shaipanich T, Hackett TL, Singhera GK, et al. ACE-2 expression in the small airway epithelia of smokers and COPD patients: implications for COVID-19. *Eur Respir J* 2020;55:2000688.
- Peters MC, Sajuthi S, Deford P, Christenson S, Rios CL, Montgomery MT, et al. COVID-19-related genes in sputum cells in asthma: relationship to demographic features and corticosteroids. *Am J Respir Crit Care Med* 2020;202:83–90.
- Streeck H, Schulte B, Kuemmerer B, Richter E, Hoeller T, Fuhrmann C. Infection fatality rate of SARS-CoV-2 infection in a German community with a super-spreading event [preprint]. medRxiv; 2020 [accessed 2019 Mar 12]. Available from: <https://www.medrxiv.org/content/10.1101/2020.05.04.20090076v2>.
- Rentsch CT, Kidwai-Khan F, Tate JP, Park LS, King JT, Skanderson M, et al. Covid-19 testing, hospital admission, and intensive care among 2,026,227 United States veterans aged 54–75 years [preprint]. medRxiv; 2020 [accessed 2019 Mar 12]. Available from: <https://www.medrxiv.org/content/10.1101/2020.04.09.20059964v1>.
- de Lusignan S, Dorward J, Correa A, Jones N, Akinyemi O, Amirthalingam G, et al. Risk factors for SARS-CoV-2 among patients in the Oxford Royal College of general practitioners research and surveillance centre primary care network: a cross-sectional study. *Lancet Infect Dis* 2020;20:1034–1042.
- Hippisley-Cox J, Young D, Coupland C, Channon KM, Tan PS, Harrison DA, et al. Risk of severe COVID-19 disease with ACE inhibitors and angiotensin receptor blockers: cohort study including 8.3 million people. *Heart* 2020;106:1503–1511.
- Halpin DMG, Faner R, Sibila O, Badia JR, Agusti A. Do chronic respiratory diseases or their treatment affect the risk of SARS-CoV-2 infection? *Lancet Respir Med* 2020;8:436–438.
- Leung JM, Niikura M, Yang CWT, Sin DD. COVID-19 and COPD. *Eur Respir J* 2020;56:2002108.
- Docherty AB, Harrison EM, Green CA, Hardwick HE, Pius R, Norman L, et al.; ISARIC4C investigators. Features of 20 133 UK patients in hospital with covid-19 using the ISARIC WHO clinical characterisation protocol: prospective observational cohort study. *BMJ* 2020;369:m1985.
- Lippi G, Henry BM. Chronic obstructive pulmonary disease is associated with severe coronavirus disease 2019 (COVID-19). *Respir Med* 2020;167:105941.
- Grasselli G, Greco M, Zanella A, Albano G, Antonelli M, Bellani G, et al.; COVID-19 Lombardy ICU Network. Risk factors associated with mortality among patients with COVID-19 in intensive care units in Lombardy, Italy. *JAMA Intern Med* 2020;180:1345–1355.
- Singh AK, Gillies CL, Singh R, Singh A, Chudasama Y, Coles B, et al. Prevalence of co-morbidities and their association with mortality in patients with COVID-19: a systematic review and meta-analysis. *Diabetes Obes Metab* [online ahead of print] 23 Jun 2020; DOI: 10.1111/dom.14124.
- Petrilli CM, Jones SA, Yang J, Rajagopalan H, O'Donnell L, Chernyak Y, et al. Factors associated with hospital admission and critical illness among 5279 people with coronavirus disease 2019 in New York City: prospective cohort study. *BMJ* 2020;369:m1966.
- Gupta S, Hayek SS, Wang W, Chan L, Mathews KS, Melamed ML, et al.; STOP-COVID Investigators. Factors associated with death in critically ill patients with coronavirus disease 2019 in the US. *JAMA Intern Med* [online ahead of print] 15 Jul 2020; DOI: 10.1001/jamainternmed.2020.3596.
- Elbeddini A, Tayefehchamani Y. Amid COVID-19 pandemic: challenges with access to care for COPD patients. *Res Social Adm Pharm* [online ahead of print] 2 Jun 2020; DOI: 10.1016/j.sapharm.2020.06.002.
- Press VG, Gershon AS, Sciruba FC, Blagev DP. Concerns about coronavirus disease-related collateral damage for patients with COPD. *Chest* 2020;158:866–868.
- Berghaus TM, Karschnia P, Haberl S, Schwaiblmair M. Disproportionate decline in admissions for exacerbated COPD during the COVID-19 pandemic. *Respir Med* 2020;10:106120.
- Chan KPF, Ma TF, Kwok WC, Leung JKC, Chiang KY, Ho JCM, et al. Significant reduction in hospital admissions for acute exacerbation of chronic obstructive pulmonary disease in Hong Kong during coronavirus disease 2019 pandemic. *Respir Med* 2020;171:106085.
- World Health Organization. Smoking and COVID-19: scientific brief. World Health Organization; 2020 [created 2020 June 30; accessed 2020 Oct 25]. Available from: <https://www.who.int/news-room/commentaries/detail/smoking-and-covid-19>.
- Fang Y, Zhang H, Xie J, Lin M, Ying L, Pang P, et al. Sensitivity of chest CT for COVID-19: comparison to RT-PCR. *Radiology* 2020;296:E115–E117.
- Yue H, Zhang M, Xing L, Wang K, Rao X, Liu H, et al. The epidemiology and clinical characteristics of co-infection of SARS-CoV-2 and influenza viruses in patients during COVID-19 outbreak. *J Med Virol* [online ahead of print] 12 Jun 2020; DOI: 10.1002/jmv.26163.
- Gousseff M, Penot P, Gallay L, Batisse D, Benech N, Bouiller K, et al.; COCOREC study group. Clinical recurrences of COVID-19 symptoms after recovery: viral relapse, reinfection or inflammatory rebound? *J Infect* 2020;81:816–846.
- Mammen MJ, Sethi S. COPD and the microbiome. *Respirology* 2016; 21:590–599.
- Khatiwada S, Subedi A. Lung microbiome and coronavirus disease 2019 (COVID-19): possible link and implications. *Hum Microb J* 2020;17:100073.
- European Respiratory Society. Recommendation from ERS Group 9.1 (Respiratory function technologists/Scientists): lung function testing during COVID-19 pandemic and beyond. European Respiratory Society; 2020 [accessed 2020 Oct 25]. Available from: <https://ers.app.box.com/s/zs1uu88wy51monr0ewd990itoz4tsn2h>.
- McCormack MC, Kaminsky DA.; American Thoracic Society. Pulmonary function laboratories: advice regarding COVID-19. American Thoracic Society; 2020 [accessed 2020 Oct 25]. Available from: <https://www.thoracic.org/professionals/clinical-resources/disease-related-resources/pulmonary-function-laboratories.php>.
- Martinez FJ, Mannino D, Leidy NK, Malley KG, Bacci ED, Barr RG, et al.; High-Risk-COPD Screening Study Group. A new approach for identifying patients with undiagnosed chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2017;195:748–756.
- Jithoo A, Enright PL, Burney P, Buist AS, Bateman ED, Tan WC, et al.; BOLD Collaborative Research Group. Case-finding options for COPD: results from the burden of obstructive lung disease study. *Eur Respir J* 2013;41:548–555.
- Mahboub B, Alzaabi A, Soriano JB, Salameh L, Mutairi YA, Yusufali AA, et al. Case-finding of chronic obstructive pulmonary disease with questionnaire, peak flow measurements and spirometry: a cross-sectional study. *BMC Res Notes* 2014;7:241.
- Perez-Padilla R, Vollmer WM, Vázquez-García JC, Enright PL, Menezes AM, Buist AS; BOLD and PLATINO Study Groups. Can a normal peak expiratory flow exclude severe chronic obstructive pulmonary disease? *Int J Tuberc Lung Dis* 2009;13:387–393.
- Aggarwal AN, Gupta D, Jindal SK. The relationship between FEV₁ and peak expiratory flow in patients with airways obstruction is poor. *Chest* 2006;130:1454–1461.
- Pothirat C, Chaiwong W, Phetsuk N, Liwsrisakun C, Bumroongkit C, Deesomchok A, et al. Peak expiratory flow rate as a surrogate for forced expiratory volume in 1 second in COPD severity classification in Thailand. *Int J Chron Obstruct Pulmon Dis* 2015;10:1213–1218.
- Llewellyn P, Sawyer G, Lewis S, Cheng S, Weatherall M, Fitzharris P, et al. The relationship between FEV₁ and PEF in the assessment of the severity of airways obstruction. *Respirology* 2002;7: 333–337.
- Jackson H, Hubbard R. Detecting chronic obstructive pulmonary disease using peak flow rate: cross sectional survey. *BMJ* 2003;327: 653–654.
- Carpenter DM, Jurdi R, Roberts CA, Hernandez M, Horne R, Chan A. A review of portable electronic spirometers: implications for asthma self-management. *Curr Allergy Asthma Rep* 2018;18:53.

39. Ramos Hernández C, Núñez Fernández M, Pallares Sanmartín A, Mouronte Roibas C, Cerdeira Domínguez L, Botana Rial MI, *et al.* Validation of the portable air-smart spirometer. *PLoS One* 2018;13:e0192789.
40. Wahidi MM, Lamb C, Murgu S, Musani A, Shojaaee S, Sachdeva A, *et al.* American Association for Bronchology and Interventional Pulmonology (AABIP) statement on the use of bronchoscopy and respiratory specimen collection in patients with suspected or confirmed COVID-19 infection. *J Bronchology Interv Pulmonol* 2020; 27:e52–e54.
41. Wahidi MM, Shojaaee S, Lamb CR, Ost D, Maldonado F, Eapen G, *et al.* The use of bronchoscopy during the coronavirus disease 2019 pandemic: CHEST/AABIP guideline and expert panel report. *Chest* 2020;158:1268–1281.
42. Wong HYF, Lam HYS, Fong AH, Leung ST, Chin TW, Lo CSY, *et al.* Frequency and distribution of chest radiographic findings in patients positive for COVID-19. *Radiology* 2020;296:E72–E78.
43. Rubin GD, Ryerson CJ, Haramati LB, Sverzellati N, Kanne JP, Raouf S, *et al.* The role of chest imaging in patient management during the COVID-19 pandemic: a multinational consensus statement from the fleischner society. *Radiology* 2020;296:172–180.
44. Rodriguez-Morales AJ, Cardona-Ospina JA, Gutiérrez-Ocampo E, Villamizar-Peña R, Holguin-Rivera Y, Escalera-Antezana JP, *et al.*; Latin American Network of Coronavirus Disease 2019-COVID-19 Research (LANCOVID-19). Clinical, laboratory and imaging features of COVID-19: a systematic review and meta-analysis. *Travel Med Infect Dis* 2020;34:101623.
45. Kulkarni S, Down B, Jha S. Point-of-care lung ultrasound in intensive care during the COVID-19 pandemic. *Clin Radiol* 2020;75:710, e1–710, e4.
46. Inui S, Fujikawa A, Jitsu M, Kunishima N, Watanabe S, Suzuki Y, *et al.* Findings in cases from the cruise ship “diamond princess” with coronavirus disease 2019 (COVID-19). *Radiol Cardiothorac Imaging* 2020;2:e200110.
47. Salehi S, Abedi A, Balakrishnan S, Gholamrezanezhad A. Coronavirus disease 2019 (COVID-19): a systematic review of imaging findings in 919 patients. *AJR Am J Roentgenol* 2020;215:87–93.
48. Wu F, Zhou Y, Wang Z, Xie M, Shi Z, Tang Z, *et al.*; Medical Treatment Expert Group for COPD and COVID-19. Clinical characteristics of COVID-19 infection in chronic obstructive pulmonary disease: a multicenter, retrospective, observational study. *J Thorac Dis* 2020; 12:1811–1823.
49. Tittaferrante S, Gupta R, Kim V; Temple University COVID-19 Research Group. Thoracic computed tomography features of coronavirus disease 2019 patients with emphysema. *Chronic Obstr Pulm Dis (Miami)* 2020;7:290–296.
50. Mossa-Basha M, Meltzer CC, Kim DC, Tuite MJ, Kolli KP, Tan BS. Radiology department preparedness for COVID-19: radiology scientific expert review panel. *Radiology* 2020;296:E106–E112.
51. Han H, Yang L, Liu R, Liu F, Wu KL, Li J, *et al.* Prominent changes in blood coagulation of patients with SARS-CoV-2 infection. *Clin Chem Lab Med* 2020;58:1116–1120.
52. Driggin E, Madhavan MV, Bikdeli B, Chuich T, Laracy J, Biondi-Zoccai G, *et al.* Cardiovascular considerations for patients, health care workers, and health systems during the COVID-19 pandemic. *J Am Coll Cardiol* 2020;75:2352–2371.
53. Guan WJ, Ni ZY, Hu Y, Liang WH, Ou CQ, He JX, *et al.*; China Medical Treatment Expert Group for Covid-19. Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med* 2020;382: 1708–1720.
54. Tang N, Bai H, Chen X, Gong J, Li D, Sun Z. Anticoagulant treatment is associated with decreased mortality in severe coronavirus disease 2019 patients with coagulopathy. *J Thromb Haemost* 2020;18: 1094–1099.
55. Llitjos JF, Leclerc M, Chochois C, Monsallier JM, Ramakers M, Auvray M, *et al.* High incidence of venous thromboembolic events in anticoagulated severe COVID-19 patients. *J Thromb Haemost* 2020; 18:1743–1746.
56. Helms J, Tacquard C, Severac F, Leonard-Lorant I, Ohana M, Delabranche X, *et al.*; CRICS TRIGGERSEP Group (Clinical Research in Intensive Care and Sepsis Trial Group for Global Evaluation and Research in Sepsis). High risk of thrombosis in patients with severe SARS-CoV-2 infection: a multicenter prospective cohort study. *Intensive Care Med* 2020;46:1089–1098.
57. Esposito S, Principi N, Leung CC, Migliori GB. Universal use of face masks for success against COVID-19: evidence and implications for prevention policies. *Eur Respir J* 2020;55:2001260.
58. Long Y, Hu T, Liu L, Chen R, Guo Q, Yang L, *et al.* Effectiveness of N95 respirators versus surgical masks against influenza: a systematic review and meta-analysis. *J Evid Based Med* 2020;13:93–101.
59. American College of Chest Physicians, American Lung Association, American Thoracic Society, COPD Foundation. Joint statement on importance of patients with chronic lung disease wearing facial coverings during COVID-19 pandemic. American College of Chest Physicians; 2020 [created 2020 July 17; accessed 2020 Oct 25]. Available from: <https://www.chestnet.org/News/Press-Releases/2020/07/Joint-Statement-on-Importance-of-Facial-Coverings>.
60. Kyung SY, Kim Y, Hwang H, Park JW, Jeong SH. Risks of N95 face mask use in subjects with COPD. *Respir Care* 2020;65:658–664.
61. Samannan R, Holt G, Calderon-Candelario R, Mirsaeidi M, Campos M. Effect of face masks on gas exchange in healthy persons and patients with COPD. *Ann Am Thorac Soc* [online ahead of print] 2 Oct 2020; DOI: 10.1513/AnnalsATS.202007-812RL.
62. Perencevich EN, Diekema DJ, Edmond MB. Moving personal protective equipment into the community: face shields and containment of COVID-19. *JAMA* 2020;323:2252–2253.
63. CDC. Considerations for wearing masks: help slow the spread of COVID-19. CDC; 2020 [accessed 2020 Oct 25]. Available from: <https://www.cdc.gov/coronavirus/2019-ncov/prevent-getting-sick/cloth-face-cover-guidance.html>.
64. Global Initiative for Chronic Obstructive Lung Disease. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease. Global Initiative for Chronic Obstructive Lung Disease; 2020 [updated 2020 Nov 4; accessed 2020 Oct 25]. Available from: <http://www.goldcopd.org/>.
65. Ergan B, Akgun M, Pacilli AMG, Nava S. Should I stay or should I go? COPD and air travel. *Eur Respir Rev* 2018;27:180030.
66. Tran K, Cimon K, Severn M, Pessoa-Silva CL, Conly J. Aerosol generating procedures and risk of transmission of acute respiratory infections to healthcare workers: a systematic review. *PLoS One* 2012;7:e35797.
67. Neufeld Z, Khataee H, Czirok A. Targeted adaptive isolation strategy for COVID-19 pandemic. *Infect Dis Model* 2020;5:357–361.
68. Butler N, Tulloch O; SSHAP. Considerations and principles for shielding people at high risk of severe outcomes from COVID-19. Institute of Developed Studies; 2020. [created 2020 May 7; accessed 2020 Oct 25]. Available from: <https://www.ids.ac.uk/publications/considerations-and-principles-for-shielding-people-at-high-risk-of-severe-outcomes-from-covid-19-april-2020>.
69. Tal-Singer R, Crapo JD. COPD at the time of COVID-19: a COPD foundation perspective. *Chronic Obstr Pulm Dis (Miami)* 2020;7: 73–75.
70. Tenforde MW, Kim SS, Lindsell CJ, Billig Rose E, Shapiro NI, Files DC, *et al.*; IVY Network Investigators; CDC COVID-19 Response Team; IVY Network Investigators. Symptom duration and risk factors for delayed return to usual health among outpatients with COVID-19 in a multistate health care systems network - United States, March-June 2020. *MMWR Morb Mortal Wkly Rep* 2020;69: 993–998.
71. Greenhalgh T, Knight M, A’Court C, Buxton M, Husain L. Management of post-acute covid-19 in primary care. *BMJ* 2020;370:m3026.
72. Carfi A, Bernabei R, Landi F; Gemelli Against COVID-19 Post-Acute Care Study Group. Persistent symptoms in patients after acute COVID-19. *JAMA* 2020;324:603–605.
73. Singanayagam A, Glanville N, Girkin JL, Ching YM, Marcellini A, Porter JD, *et al.* Corticosteroid suppression of antiviral immunity increases bacterial loads and mucus production in COPD exacerbations. *Nat Commun* 2018;9:2229.
74. Skevaki CL, Christodoulou I, Spyridaki IS, Tiniakou I, Georgiou V, Xepapadaki P, *et al.* Budesonide and formoterol inhibit inflammatory mediator production by bronchial epithelial cells infected with rhinovirus. *Clin Exp Allergy* 2009;39:1700–1710.

75. Thomas BJ, Porritt RA, Hertzog PJ, Bardin PG, Tate MD. Glucocorticosteroids enhance replication of respiratory viruses: effect of adjuvant interferon. *Sci Rep* 2014;4:7176.
76. Yamaya M, Nishimura H, Deng X, Sugawara M, Watanabe O, Nomura K, et al. Inhibitory effects of glycopyrronium, formoterol, and budesonide on coronavirus HCoV-229E replication and cytokine production by primary cultures of human nasal and tracheal epithelial cells. *Respir Investig* 2020;58:155–168.
77. Halpin DMG, Singh D, Hadfield RM. Inhaled corticosteroids and COVID-19: a systematic review and clinical perspective. *Eur Respir J* 2020;55:2001009.
78. Schultze A, Walker AJ, MacKenna B, Morton CE, Bhaskaran K, Brown JP, et al. Inhaled corticosteroid use and risk COVID-19 related death among 966,461 patients with COPD or asthma: an OpenSAFELY analysis [preprint]. *MedRxiv*; 2020 [created 2020 June 20; accessed 2020 Jun 20] Available from: <https://www.medrxiv.org/content/10.1101/2020.06.19.20135491v1>.
79. Singh D, Halpin DMG. Inhaled corticosteroids and COVID-19-related mortality: confounding or clarifying? *Lancet Respir Med* 2020;8:1065–1066.
80. O'Neil CA, Li J, Leavey A, Wang Y, Hink M, Wallace M, et al.; Centers for Disease Control and Prevention Epicenters Program. Characterization of aerosols generated during patient care activities. *Clin Infect Dis* 2017;65:1335–1341.
81. Simonds AK, Hanak A, Chatwin M, Morrell M, Hall A, Parker KH, et al. Evaluation of droplet dispersion during non-invasive ventilation, oxygen therapy, nebuliser treatment and chest physiotherapy in clinical practice: implications for management of pandemic influenza and other airborne infections. *Health Technol Assess* 2010;14:131–172.
82. van Doremalen N, Bushmaker T, Morris DH, Holbrook MG, Gamble A, Williamson BN, et al. Aerosol and surface stability of SARS-CoV-2 as compared with SARS-CoV-1. *N Engl J Med* 2020;382:1564–1567.
83. Heinzerling A, Stuckey MJ, Scheuer T, Xu K, Perkins KM, Resseger H, et al. Transmission of COVID-19 to health care personnel during exposures to a hospitalized patient - Solano County, California, February 2020. *MMWR Morb Mortal Wkly Rep* 2020;69:472–476.
84. Tashkin DP, Barjaktarevic IZ. Nebulized treatments and the possible risk of coronavirus transmission: where is the evidence? *Chronic Obstr Pulm Dis (Miami)* 2020;7:136–138.
85. Respiratory Care Committee of Chinese Thoracic Society. Expert consensus on preventing nosocomial transmission during respiratory care for critically ill patients infected by 2019 novel coronavirus pneumonia [in Chinese]. *Zhonghua Jie He He Hu Xi Za Zhi* 2020;43:288–296.
86. Salisbury H. Helen Salisbury: how will we run flu clinics in a pandemic? *BMJ* 2020;370:m3033.
87. Demeyer H, Louvaris Z, Frei A, Rabinovich RA, de Jong C, Gimeno-Santos E, et al.; Mr Papp PROactive study group and the PROactive consortium. Physical activity is increased by a 12-week semiautomated telecoaching programme in patients with COPD: a multicentre randomised controlled trial. *Thorax* 2017;72:415–423.
88. American Thoracic Society. Assembly on pulmonary rehabilitation: guidance for re-opening-opening pulmonary rehabilitation programs. American Thoracic Society; 2020 [accessed 2020 Oct 25]. Available from: <https://www.thoracic.org/members/assemblies/assemblies/pr/resources/ats-pr-assembly-re-opening-pr-document-final.pdf>.
89. Nield M, Hoo GW. Real-time telehealth for COPD self-management using Skype™. *COPD* 2012;9:611–619.
90. Beigel JH, Tomashek KM, Dodd LE, Mehta AK, Zingman BS, Kalil AC, et al.; ACTT-1 Study Group Members. Remdesivir for the treatment of covid-19 - final report. *N Engl J Med* [online ahead of print] 8 Oct 2020; DOI: 10.1056/NEJMoa2007764.
91. Horby P, Lim WS, Emberson JR, Mafham M, Bell JL, Linsell L, et al.; RECOVERY Collaborative Group. Dexamethasone in hospitalized patients with covid-19 - preliminary report. *N Engl J Med* [online ahead of print] 17 Jul 2020; DOI: 10.1056/NEJMoa2021436.
92. Muhammad S, Long X, Salman M. COVID-19 pandemic and environmental pollution: a blessing in disguise? *Sci Total Environ* 2020;728:138820.
93. Hewitt R, Farne H, Ritchie A, Luke E, Johnston SL, Mallia P. The role of viral infections in exacerbations of chronic obstructive pulmonary disease and asthma. *Thorax* 2016;10:158–174.
94. Ackermann M, Verleden SE, Kuehnel M, Haverich A, Welte T, Laenger F, et al. Pulmonary vascular endothelialitis, thrombosis, and angiogenesis in covid-19. *N Engl J Med* 2020;383:120–128.
95. Calabrese F, Pezzuto F, Fortarezza F, Hofman P, Kern I, Panizo A, et al. Pulmonary pathology and COVID-19: lessons from autopsy. The experience of European Pulmonary Pathologists. *Virchows Arch* 2020;477:359–372.
96. Wedzicha JA, Singh R, Mackay AJ. Acute COPD exacerbations. *Clin Chest Med* 2014;35:157–163.
97. Argenziano MG, Bruce SL, Slater CL, Tiao JR, Baldwin MR, Barr RG, et al. Characterization and clinical course of 1000 patients with coronavirus disease 2019 in New York: retrospective case series. *BMJ* 2020;369:m1996.
98. Malik P, Patel U, Mehta D, Patel N, Kelkar R, Akrmah M, et al. Biomarkers and outcomes of COVID-19 hospitalisations: systematic review and meta-analysis. *BMJ Evid Based Med* [online ahead of print] 15 Sep 2020; DOI: 10.1136/bmjebm-2020-111536.
99. Shang L, Zhao J, Hu Y, Du R, Cao B. On the use of corticosteroids for 2019-nCoV pneumonia. *Lancet* 2020;395:683–684.
100. Dagens A, Sigfrid L, Cai E, Lipworth S, Cheng V, Harris E, et al. Scope, quality, and inclusivity of clinical guidelines produced early in the covid-19 pandemic: rapid review. *BMJ* 2020;369:m1936.
101. Arabi YM, Mandourah Y, Al-Hameed F, Sindi AA, Almekhlafi GA, Hussein MA, et al.; Saudi Critical Care Trial Group. Corticosteroid therapy for critically ill patients with Middle East respiratory syndrome. *Am J Respir Crit Care Med* 2018;197:757–767.
102. Lee N, Allen Chan KC, Hui DS, Ng EK, Wu A, Chiu RW, et al. Effects of early corticosteroid treatment on plasma SARS-associated Coronavirus RNA concentrations in adult patients. *J Clin Virol* 2004;31:304–309.
103. Lee N, Chan PK, Hui DS, Rainer TH, Wong E, Choi KW, et al. Viral loads and duration of viral shedding in adult patients hospitalized with influenza. *J Infect Dis* 2009;200:492–500.
104. Russell CD, Millar JE, Baillie JK. Clinical evidence does not support corticosteroid treatment for 2019-nCoV lung injury. *Lancet* 2020;395:473–475.
105. World Health Organization. Clinical management of COVID-19: interim guidance. World Health Organization; 2020 [created 2020 May 27; accessed 2020 Oct 25]. Available from: <https://www.who.int/publications/i/item/clinical-management-of-covid-19>.
106. Wu C, Chen X, Cai Y, Xia J, Zhou X, Xu S, et al. Risk factors associated with acute respiratory distress syndrome and death in patients with coronavirus disease 2019 pneumonia in Wuhan, China. *JAMA Intern Med* 2020;180:934–943.
107. Sterne JAC, Murthy S, Diaz JV, Slutsky AS, Villar J, Angus DC, et al.; WHO Rapid Evidence Appraisal for COVID-19 Therapies (REACT) Working Group. Association between administration of systemic corticosteroids and mortality among critically ill patients with COVID-19: a meta-analysis. *JAMA* 2020;324:1330–1341.
108. Rawson TM, Moore LSP, Zhu N, Ranganathan N, Skolimowska K, Gilchrist M, et al. Bacterial and fungal co-infection in individuals with coronavirus: a rapid review to support COVID-19 antimicrobial prescribing. *Clin Infect Dis* [Online ahead of print] 2 May 2020; DOI: 10.1093/cid/ciaa530.
109. Verroken A, Scohy A, Gérard L, Wittebole X, Collienne C, Laterre PF. Co-infections in COVID-19 critically ill and antibiotic management: a prospective cohort analysis. *Crit Care* 2020;24:410.
110. Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 2020;395:1054–1062.
111. Jiang DH, McCoy RG. Planning for the post-COVID syndrome: how payers can mitigate long-term complications of the pandemic. *J Gen Intern Med* 2020;35:3036–3039.
112. Borczuk AC, Salvatore SP, Seshan SV, Patel SS, Bussel JB, Mostyka M, et al. COVID-19 pulmonary pathology: a multi-institutional autopsy cohort from Italy and New York City. *Modern Pathology* 2020;33:2156–2168.

113. Gattinoni L, Chiumello D, Rossi S. COVID-19 pneumonia: ARDS or not? *Crit Care* 2020;24:154.
114. Gattinoni L, Coppola S, Cressoni M, Busana M, Rossi S, Chiumello D. COVID-19 does not lead to a “typical” acute respiratory distress syndrome. *Am J Respir Crit Care Med* 2020;201:1299–1300.
115. Panwar R, Madotto F, Laffey JG, Van Haren FMP. Compliance phenotypes in early acute respiratory distress syndrome before the COVID-19 pandemic. *Am J Respir Crit Care Med* 2020;202:1244–1252.
116. Brault C, Zerbib Y, Kontar L, Fouquet U, Carpentier M, Metzler M, et al. COVID-19– versus non-COVID-19–related acute respiratory distress syndrome: differences and similarities. *Am J Respir Crit Care Med* 2020;202:1301–1304.
117. Grieco DL, Bongiovanni F, Chen L, Menga LS, Cutuli SL, Pintaudi G, et al. Respiratory physiology of COVID-19-induced respiratory failure compared to ARDS of other etiologies. *Crit Care* 2020;24:529.
118. Lechowicz K, Drożdżał S, Machaj F, Rosik J, Szostak B, Zegan-Barańska M, et al. COVID-19: the potential treatment of pulmonary fibrosis associated with SARS-CoV-2 infection. *J Clin Med* 2020;9:1917.
119. Rimmelink M, De Mendonça R, D’Haene N, De Clercq S, Verocq C, Lebrun L, et al. Unspecific post-mortem findings despite multiorgan viral spread in COVID-19 patients. *Crit Care* 2020;24:495.
120. Palmer K, Monaco A, Kivipelto M, Onder G, Maggi S, Michel JP, et al. The potential long-term impact of the COVID-19 outbreak on patients with non-communicable diseases in Europe: consequences for healthy aging. *Aging Clin Exp Res* 2020;32:1189–1194.
121. Zaim S, Chong JH, Sankaranarayanan V, Harky A. COVID-19 and multiorgan response. *Curr Probl Cardiol* 2020;45:100618.
122. Puelles VG, Lütgehetmann M, Lindenmeyer MT, Sperhake JP, Wong MN, Allweiss L, et al. Multiorgan and renal tropism of SARS-CoV-2. *N Engl J Med* 2020;383:590–592.
123. Dobesh PP, Trujillo TC. Coagulopathy, venous thromboembolism, and anticoagulation in patients with COVID-19. *Pharmacotherapy* [online ahead of print] 1 Oct 2020; DOI: 10.1002/phar.2465.
124. Ambrosetti M, Ageno W, Spanevello A, Salerno M, Pedretti RF. Prevalence and prevention of venous thromboembolism in patients with acute exacerbations of COPD. *Thromb Res* 2003;112:203–207.
125. Kim V, Goel N, Gangar J, Zhao H, Ciccolella DE, Silverman EK, et al.; COPD Gene Investigators. Risk factors for venous thromboembolism in chronic obstructive pulmonary disease. *Chronic Obstr Pulm Dis (Miami)* 2014;1:239–249.
126. Paranjpe I, Fuster V, Lala A, Russak AJ, Glicksberg BS, Levin MA, et al. Association of treatment dose anticoagulation with in-hospital survival among hospitalized patients with COVID-19. *J Am Coll Cardiol* 2020;76:122–124.
127. Wu Z, McGoogan JM. Characteristics of and important lessons from the coronavirus disease 2019 (COVID-19) outbreak in China: summary of a report of 72314 cases from the Chinese center for disease control and prevention. *JAMA* 2020;323:1239–1242.
128. Qiu H, Tong Z, Ma P, Hu M, Peng Z, Wu W, et al.; China Critical Care Clinical Trials Group. Intensive care during the coronavirus epidemic. *Intensive Care Med* 2020;46:576–578.
129. Johns Hopkins University & Medicine. Johns Hopkins Coronavirus Resource Center. Johns Hopkins University & Medicine; 2020 [accessed 2020 Oct 25]. Available from: <https://coronavirus.jhu.edu>.
130. Schünemann HJ, Khabsa J, Solo K, Khamsi AM, Brignardello-Petersen R, El-Harakeh A, et al. Ventilation techniques and risk for transmission of coronavirus disease, including COVID-19: a living systematic review of multiple streams of evidence. *Ann Intern Med* 2020;173:204–216.
131. Grasselli G, Zangrillo A, Zanella A, Antonelli M, Cabrini L, Castelli A, et al.; COVID-19 Lombardy ICU Network. Baseline characteristics and outcomes of 1591 patients infected with SARS-CoV-2 admitted to ICUs of the Lombardy region, Italy. *JAMA* 2020;323:1574–1581.
132. Namendys-Silva SA. Respiratory support for patients with COVID-19 infection. *Lancet Respir Med* 2020;8:e18.
133. Kluge S, Janssens U, Welte T, Weber-Carstens S, Marx G, Karagiannidis C. German recommendations for critically ill patients with COVID-19. *Med Klin Intensivmed Notf Med* [online ahead of print] 14 Apr 2020; DOI: 10.1007/s00063-020-00689-w.
134. Cheung JC, Ho LT, Cheng JW, Cham EYK, Lam KN. Staff safety during emergency airway management for COVID-19 in Hong Kong. *Lancet Respir Med* 2020;8:e19.
135. Crimi C, Noto A, Cortegiani A, Impellizzeri P, Elliott M, Ambrosino N, et al. Noninvasive respiratory support in acute hypoxemic respiratory failure associated with COVID-19 and other viral infections. *Minerva Anesthesiol* [online ahead of print] 5 Aug 2020; DOI: 10.23736/S0375-9393.20.14785-0.
136. Patel M, Gangemi A, Marron R, Chowdhury J, Yousef I, Zheng M, et al. Retrospective analysis of high flow nasal therapy in COVID-19-related moderate-to-severe hypoxaemic respiratory failure. *BMJ Open Respir Res* 2020;7:e000650.
137. Demoule A, Vieillard Baron A, Darmon M, Beurton A, Géri G, Voiriot G, et al. High-flow nasal cannula in critically ill patients with severe COVID-19. *Am J Respir Crit Care Med* 2020;202:1039–1042.
138. Frat JP, Thille AW, Mercat A, Girault C, Ragot S, Perbet S, et al.; FLORALI Study Group; REVA Network. High-flow oxygen through nasal cannula in acute hypoxemic respiratory failure. *N Engl J Med* 2015;372:2185–2196.
139. Ni YN, Luo J, Yu H, Liu D, Liang BM, Liang ZA. The effect of high-flow nasal cannula in reducing the mortality and the rate of endotracheal intubation when used before mechanical ventilation compared with conventional oxygen therapy and noninvasive positive pressure ventilation: a systematic review and meta-analysis. *Am J Emerg Med* 2018;36:226–233.
140. Alhazzani W, Möller MH, Arabi YM, Loeb M, Gong MN, Fan E, et al. Surviving sepsis campaign: guidelines on the management of critically ill adults with coronavirus disease 2019 (COVID-19). *Crit Care Med* 2020;48:e440–e469.
141. Telias I, Katira BH, Brochard L. Is the prone position helpful during spontaneous breathing in patients with COVID-19? *JAMA* 2020;323:2265–2267.
142. Slutsky AS, Ranieri VM. Ventilator-induced lung injury. *N Engl J Med* 2013;369:2126–2136.
143. Berlin DA, Gulick RM, Martinez FJ. Severe covid-19. *N Engl J Med* [online ahead of print] 15 May 2020; DOI: 10.1056/NEJMc2009575.
144. Fan E, Beitler JR, Brochard L, Calfee CS, Ferguson ND, Slutsky AS, et al. COVID-19-associated acute respiratory distress syndrome: is a different approach to management warranted? *Lancet Respir Med* 2020;8:816–821.
145. Bellani G, Laffey JG, Pham T, Madotto F, Fan E, Brochard L, et al.; LUNG SAFE Investigators; ESICM Trials Group. Noninvasive ventilation of patients with acute respiratory distress syndrome: insights from the LUNG SAFE study. *Am J Respir Crit Care Med* 2017;195:67–77.
146. Ma X, Liang M, Ding M, Liu W, Ma H, Zhou X, et al. Extracorporeal membrane oxygenation (ECMO) in critically ill patients with coronavirus disease 2019 (COVID-19) pneumonia and acute respiratory distress syndrome (ARDS). *Med Sci Monit* 2020;26:e925364.
147. Bartlett RH, Ogino MT, Brodie D, McMullan DM, Lorusso R, MacLaren G, et al. Initial ELSO guidance document: ECMO for COVID-19 patients with severe cardiopulmonary failure. *ASAIO J* 2020;66:472–474.
148. MacLaren G, Fisher D, Brodie D. Preparing for the most critically ill patients with COVID-19: the potential role of extracorporeal membrane oxygenation. *JAMA* 2020;323:1245–1246.
149. Ramanathan K, Antognini D, Combes A, Paden M, Zakhary B, Ogino M, et al. Planning and provision of ECMO services for severe ARDS during the COVID-19 pandemic and other outbreaks of emerging infectious diseases. *Lancet Respir Med* 2020;8:518–526.
150. Shekar K, Badulak J, Peek G, Boeken U, Dalton HJ, Arora L, et al.; ELSO Guideline Working Group*. Extracorporeal life support organization COVID-19 interim guidelines. *ASAIO J* [online ahead of print] 29 Apr 2020; DOI: 10.1097/MAT.0000000000001193.

151. Hamele M, Neumayer K, Sweney J, Poss WB. Always ready, always prepared—preparing for the next pandemic. *Transl Pediatr* 2018;7: 344–355.
152. Zochios V, Brodie D, Charlesworth M, Parhar KK. Delivering extracorporeal membrane oxygenation for patients with COVID-19: what, who, when and how? *Anaesthesia* 2020;75:997–1001.
153. Li J, Fink JB, Ehrmann S. High-flow nasal cannula for COVID-19 patients: low risk of bio-aerosol dispersion. *Eur Respir J* 2020;55: 2000892.
154. Raboud J, Shigayeva A, McGeer A, Bontovics E, Chapman M, Gravel D, et al. Risk factors for SARS transmission from patients requiring intubation: a multicentre investigation in Toronto, Canada. *PLoS One* 2010;5:e10717.
155. Hautmann H, Gamarra F, Pfeifer KJ, Huber RM. Fiberoptic bronchoscopic balloon dilatation in malignant tracheobronchial disease: indications and results. *Chest* 2001;120:43–49.
156. Pfeifer M, Ewig S, Voshaar T, Randerath WJ, Bauer T, Geiseler J, et al. Position paper for the state-of-the-art application of respiratory support in patients with COVID-19. *Respiration* 2020;99:521–542.
157. Shaw KM, Lang AL, Lozano R, Szabo M, Smith S, Wang J. Intensive care unit isolation hood decreases risk of aerosolization during noninvasive ventilation with COVID-19. *Can J Anaesth* 2020;67: 1481–1483.
158. Schols AM, Broekhuizen R, Weling-Scheepers CA, Wouters EF. Body composition and mortality in chronic obstructive pulmonary disease. *Am J Clin Nutr* 2005;82:53–59.
159. Needham DM, Davidson J, Cohen H, Hopkins RO, Weinert C, Wunsch H, et al. Improving long-term outcomes after discharge from intensive care unit: report from a stakeholders' conference. *Crit Care Med* 2012;40:502–509.
160. Needham DM, Feldman DR, Kho ME. The functional costs of ICU survivorship: collaborating to improve post-ICU disability. *Am J Respir Crit Care Med* 2011;183:962–964.
161. Herridge MS, Chu LM, Matte A, Tomlinson G, Chan L, Thomas C, et al.; RECOVER Program Investigators (Phase 1: towards RECOVER); Canadian Critical Care Trials Group. The RECOVER program: disability risk groups and 1-year outcome after 7 or more days of mechanical ventilation. *Am J Respir Crit Care Med* 2016; 194:831–844.
162. Griffith DM, Salisbury LG, Lee RJ, Lone N, Merriweather JL, Walsh TS; RECOVER Investigators. Determinants of health-related quality of life after ICU: importance of patient demographics, previous comorbidity, and severity of illness. *Crit Care Med* 2018;46: 594–601.
163. Holm SE, Mu K. Discharge planning for the elderly in acute care: the perceptions of experienced occupational therapists. *Phys Occup Ther Geriatr* 2012;30:214–228.
164. Spruit MA, Holland AE, Singh SJ, Tonia T, Wilson KC, Troosters T. COVID-19: interim guidance on rehabilitation in the hospital and post-hospital phase from a European Respiratory Society and American Thoracic Society-coordinated international task force. *Eur Respir J* 2020;10: 2002197.
165. Hui DS, Joynt GM, Wong KT, Gomersall CD, Li TS, Antonio G, et al. Impact of severe acute respiratory syndrome (SARS) on pulmonary function, functional capacity and quality of life in a cohort of survivors. *Thorax* 2005;60:401–409.
166. Das KM, Lee EY, Singh R, Enani MA, Al Dossari K, Van Gorkom K, et al. Follow-up chest radiographic findings in patients with MERS-CoV after recovery. *Indian J Radiol Imaging* 2017;27:342–349.
167. Gandhi RT, Lynch JB, Del Rio C. Mild or moderate covid-19. *N Engl J Med* 2020;383:1757–1766.
168. Alqahtani JS, Oyelade T, Aldhahir AM, Alghamdi SM, Almeahmadi M, Alqahtani AS, et al. Prevalence, severity and mortality associated with COPD and smoking in patients with COVID-19: a rapid systematic review and meta-analysis. *PLoS One* 2020;15:e0233147.
169. Hosey MM, Needham DM. Survivorship after COVID-19 ICU stay. *Nat Rev Dis Primers* 2020;6:60.
170. Lamas D. Chronic critical illness. *N Engl J Med* 2014;370:175–177.
171. Tracy CS, Bell SH, Nickell LA, Charles J, Upshur RE. The IMPACT clinic: innovative model of interprofessional primary care for elderly patients with complex health care needs. *Can Fam Physician* 2013; 59:e148–e155.